

11. FEJEZET. ENDOKRIN EREDETŰ BELGYÓGYÁSZATI KÓRFORMÁK

ÁLTALÁNOS ISMERETEK

Beatrice, L. és mtsai (2018): Concurrent endocrine neoplasias in dogs and cats: a retrospective study (2004-2014). *Vet. Rec.* 182, 323. doi: 10.1136/vr.104199.

McLean, J.L. és mtsai (2014): Worldwide prevalence and risk factors for feline hyperthyroidism: A review. *J. S. Afr. Vet. Assoc.* 85, 1097. doi: 10.4102/jsava.v85i1.1097.

O'Neill, D.G. és mtsai (2016): Epidemiology of diabetes mellitus among 193,435 cats attending primary-care veterinary practices in England. *J. Vet. Intern. Med.* 30, 964-972.

Pappa, T. és mtsai (2015): Inherited defects of thyroxine-binding proteins. *Best Pract. Res. Clin. Endocrinol. Metab.* 29, 735-747.

Salesov, E. és mtsai (2015): Urinary and plasma catecholamines and metanephrines in dogs with pheochromocytoma, hypercortisolism, nonadrenal disease and in healthy dogs. *J. Vet. Intern. Med.* 29, 597-602.

ENDOKRIN BETEGSÉGEK

Altschul, M. és mtsai (1997): Evaluation of somatostatin analogues for the detection and treatment of gastrinoma in a dog. *J. Small Anim. Pract.* 38, 286-291.

Asakawa, M.G. és mtsai (2013): Necrolytic migratory erythema associated with a glucagon-producing primary hepatic neuroendocrine carcinoma in a cat. *Vet. Dermatol.* 24, 466– 469.

Ash, R.A. és mtsai (2005): Primary hyperaldosteronism in the cat: a series of 13 cases. *J. Feline Med. Surg.* 7, 173-182.

Barber, P.J. (2004): Disorders of the parathyroid glands. *J. Feline Med. Surg.* 6, 259-269.

Bell, E.T. és mtsai (2012): Immune-mediated myasthenia gravis in a methimazole-treated cat. *J. Small Anim. Pract.* 53, 661-663.

Bertolini, G. és mtsai (2017): Incidental and nonincidental canine thyroid tumors assessed by multidetector row computed tomography: a single-centre cross sectional study in 4520 dogs. *Vet. Radiol. Ultrasound.* 58, 304-314.

Buishand, F.O. és mtsai (2018): Utility of contrast-enhanced computed tomography in the evaluation of canine insulinoma location. *Vet. Q.* 38, 53-62.

Cartwright, J.A. (2016): Polyglandular endocrinopathy type II (Schmidt's syndrome) in a Dobermann pinscher. *J. Small Anim. Pract.* 57, 491-494.

Csöndes, J., Fábrián, I. Szabó, B., Máthé, Á., Vajdovich, P. (2017): Assessment of adrenocortical reserve capacity and inflammatory parameters in critically ill dogs. *Acta Vet. Hung.* 65, 475-486.

- Fabres, V. és mtsai (2018): Evaluation of telmisartan administration as a suppression test for primary hyperaldosteronism diagnosis in cats. Research communications of the 28th ECVIM-CA Congress. <https://doi.org/10.1111/jvim.15372>
- Feldman, E.C. és mtsai (2005): Pretreatment clinical and laboratory findings in dogs with primary hyperparathyroidism: 210 cases (1987-2004). *J. Am. Vet. Med. Assoc.* 227, 756-761.
- Galac, S. és mtsai (2010): Expression of receptors for luteinizing hormone, gastric-inhibitory polypeptide, and vasopressin in normal adrenal glands and cortisol-secreting adrenocortical tumors in dogs. *Domest. Anim. Endocrinol.* 39, 63-75.
- Giunti, M. és mtsai (2017): Retrospective evaluation of circulating thyroid hormones in critically ill dogs with systemic inflammatory response syndrome. *J. Vet. Sci.* 18, 471-477.
- Goemans, A.F. és mtsai (2017): Validation and determination of a reference interval for canine HbA1c using an immunoturbidimetric assay. *Vet. Clin. Pathol.* 46, 227-237.
- Gostelow, R. és mtsai (2017): Pasireotide long-acting release treatment for diabetic cats with underlying hypersomatotropism. *J. Vet. Intern. Med.* 31, 355-364.
- Graham, K.J. és mtsai (2012): Intraoperative parathyroid hormone concentration to confirm removal of hypersecretory parathyroid tissue and time to postoperative normocalcaemia in nine dogs with primary hyperparathyroidism. *Aust. Vet. J.* 90, 203-209.
- Greene, S.N. és Bright, R.M. (2008): Insulinoma in a cat. *J. Small Anim. Pract.* 49, 38-40.
- Gunn, E. és mtsai (2016): Hydrocortisone in the management of acute hypoadrenocorticism in dogs: a retrospective series of 30 cases. *J. Small Anim. Pract.* 57, 227-233.
- Guttin, T. és mtsai (2015): Outcomes for dogs with primary hyperparathyroidism following treatment with percutaneous ultrasound-guided ethanol ablation of presumed functional parathyroid nodules: 27 cases (2008-2011). *J. Am. Vet. Med. Assoc.* 247, 771-777.
- Ham, K. és mtsai (2009): Validation of a rapid parathyroid hormone assay and measurement of parathyroid hormone in dogs with naturally occurring primary hyperparathyroidism. *Vet. Surg.* 38, 122-132.
- Johnson, K.D. és mtsai (2006): Primary hyperaldosteronism in a dog with concurrent lymphoma. *J. Vet. Med. A. Physiol. Pathol. Clin. Med.* 53, 467-470.
- Johnstone, T. és mtsai (2014): Hypothyroidism associated with acromegaly and insulin-resistant diabetes mellitus in a Samoyed. *Aust. Vet. J.* 92, 437-442.
- Johnstone, T. és mtsai (2014): Hypothyroidism associated with acromegaly and insulin-resistant diabetes mellitus in a Samoyed. *Aust. Vet. J.* 92, 437-442.
- Kippenes, H. és mtsai (2001): Mensuration of the normal pituitary gland from magnetic resonance images in 96 dogs. *Vet. Radiol. Ultrasound.* 42, 130-133.

- Kirsten, L. (2019): Concurrent pituitary and adrenocortical lesions on computed tomography imaging in dogs with spontaneous hypercortisolism. *J. Vet. Intern. Med.* 33, 72–78.
- Knottenbelt és Herrtage (2002): Use of proligestone in the management of three German shepherd dogs with pituitary dwarfism. *J. Small Anim. Pract.* 43. 164-170.
- Kooistra, H.S. és mtsai (2000): Combined pituitary hormone deficiency in German Shepherd dogs with dwarfism. *Domest. Anim. Endocrinol.* 19, 177-190.
- Lathan, P. és Thompson, A.L. (2018): Management of hypoadrenocorticism (Addison's disease) in dogs. *Vet. Med. (Auckl).* 9, 1-10.
- Lautenschlaeger, I.E. és mtsai (2013): Comparison between computed tomography and (99m)TC-pertechnetate scintigraphy characteristics of the thyroid gland in cats with hyperthyroidism. *Vet. Radiol. Ultrasound.* 54, 666-673.
- Lehner, L., Czeibert, K., Csöndes, J., Balogh, N., Kerekes, Z., Jakab, C. (2018): Az agyalapi mirigy daganatának részleges eltávolítása endoszkóp segítségével az ékcsonton keresztül kutyában. *Esetismertetés. Magy. Állatorvosok Lapja* 140, 535-550.
- Long, C.D. és mtsai (1999): Percutaneous ultrasound-guided chemical parathyroid ablation for treatment of primary hyperparathyroidism in dogs. *J. Am. Vet. Med. Assoc.* 215, 217-221.
- Lurye, J.C. és Behrend, E.N. (2001): Endocrine tumors. *Vet. Clin. North Am. Small Anim. Pract.* 3, 1083-1110.
- MacDonald, K.A. és mtsai (2008): Effect of spironolactone on diastolic function and left ventricular mass in Maine Coon cats with familial hypertrophic cardiomyopathy. *J. Vet. Intern. Med.* 22, 335-341.
- Mayer, M.N. és Treuil, P.L. (2007): Radiation therapy for pituitary tumors in the dog and cat. *Can. Vet. J.* 48, 316-318.
- Meij, B.P. (2001): Hypophysectomy as a treatment for canine and feline Cushing's disease. *Vet. Clin. North Am. Small Anim. Pract.* 31, 1015-1041.
- Meij, B.P. (2001): Transsphenoidal hypophysectomy for treatment of pituitary-dependent hyperadrenocorticism in 7 cats. *Vet. Surg.* 30, 72-86.
- Merryman, J.I. és mtsai (1999): Overexpression of c-ras in hyperplasia and adenomas of the feline thyroid gland: an immunohistochemical analysis of 34 cases. *Vet Pathol.* 36, 117.
- Miller, M.L. és mtsai (2017): Thyroid cysts in cats: A retrospective study of 40 cases. *J. Vet. Intern. Med.* 31, 723-729.
- Morgan M.J. és mtsai (2008): Clinical peripheral neuropathy associated with diabetes mellitus in 3 dogs. *Can. Vet. J.* 49, 583-586.
- Neiger, R. és mtsai (2004): Trilostane therapy for treatment of pituitary-dependent hyperadrenocorticism in 5 cats. *J. Vet. Intern. Med.* 18, 160-164.

- Niessen, S. és mtsai (2014): Pasireotide (SOM230) opens doors to medical management of feline hypersomatotropism. *J. Vet. Intern. Med.* 27, 685.
- Oberkirchner, U. és mtsai (2010): Successful treatment of canine necrolytic migratory erythema (superficial necrolytic dermatitis) due to metastatic glucagonoma with octreotide. *Vet. Dermatol.* 21, 510–516.
- Pagani, E. és mtsai (2016): Ultrasonographic features of adrenal gland lesions in dogs can aid in diagnosis. *BMC Vet. Res.* 12, 267.
- Peterson, M.E. (2006): Radioiodine treatment of hyperthyroidism. *Clin. Tech. Small Anim. Pract.* 21, 34-39.
- Peterson, M.E. és Broome, M.R. (2015): Thyroid scintigraphy findings in 2096 cats with hyperthyroidism. *Vet. Radiol. Ultrasound.* 56, 84-95.
- Peterson, M.E. és mtsai (2018): Spontaneous primary hypothyroidism in 7 adult cats. *J. Vet. Intern. Med.* 32, 1864-1873.
- Pijnacker, T. (2018): Use of basal and TRH-stimulated plasma growth hormone concentrations to differentiate between primary hypothyroidism and nonthyroidal illness in dogs. *J. Vet. Intern. Med.* 32, 1319-1324.
- Pijnacker, T. és mtsai (2018): Use of basal and TRH-stimulated plasma growth hormone concentrations to differentiate between primary hypothyroidism and nonthyroidal illness in dogs. *J. Vet. Intern. Med.* 32, 1319-1324.
- Pollard, R.E. és mtsai (2001): Percutaneous ultrasonographically guided radiofrequency heat ablation for treatment of primary hyperparathyroidism in dogs. *J. Am. Vet. Med. Assoc.* 218, 1106-1110.
- Rasor, L. és mtsai (2007): Retrospective evaluation of three treatment methods for primary hyperparathyroidism in dogs. *J. Am. Anim. Hosp. Assoc.* 43, 70-77.
- Russell, N.J. és mtsai (2006): Primary hypoparathyroidism in dogs: a retrospective study of 17 cases. *Aust. Vet. J.* 84, 285-290.
- Sanders, K. és mtsai (2018): Treating canine Cushing's syndrome: Current options and future prospects. *Vet. J.* 241, 42-51.
- Sawyer, E.S. és mtsai (2011): Outcome of 19 dogs with parathyroid carcinoma after surgical excision. *Vet. Comp. Oncol.* 10, 57-64.
- Schaub, S. és Wigger, A. (2013): Ultrasound-aided diagnosis of an insulinoma in a cat. *Tierärztl. Prax. Ausg. K. Kleintiere Heimtiere.* 41, 338-342.
- Taylor, J.A. és mtsai (2014): Functional ectopic adrenal carcinoma in a dog. *Can. Vet. J.* 55, 845-848.

Van Bokhorst, K.L: és mtsai (2019): Concurrent pituitary and adrenocortical lesions on computed

Van der Vlugt-Meijer, R.H. és mtsai (2003): Dynamic computed tomography of the pituitary gland in dogs with pituitary-dependent hyperadrenocorticism. *J. Vet. Intern. Med.* 17, 773-780.

Van Vonderen, I.K. és mtsai (2004): Vasopressin response to osmotic stimulation in 18 young dogs with polyuria and polydipsia. *J. Vet. Intern. Med.* 18, 800-806.

Wakayama, J.A. és mtsai (2017): A retrospective study of dogs with atypical hypoadrenocorticism: a diagnostic cut-off or continuum? *J. Small Anim. Pract.* 58,365-371.

Walczak, R. és mtsai (2019): Canine insulinomas appear hyperintense on MRI T2-weighted images and isointense on T1-weighted images. *Vet. Radiol. Ultrasound.* 60, 330-337.

Ward, C.R. (2007): Feline thyroid storm. *Vet. Clin. North Am. Small Anim. Pract.* 37, 745-754.

Ward, C.R. és mtsai (2010): Evaluation of activation of G proteins in response to thyroid stimulating hormone in thyroid gland cells from euthyroid and hyperthyroid cats. *Am J Vet Res.* 71, 643-648.