

4. FEJEZET. A MÁJ ÉS AZ EPEUTAK BETEGSÉGEI

KÓRHATÁROZÁS

Bahr, K.L. és mtsai (2013): Accuracy of US-guided FNA of focal liver lesions in dogs: 140 cases (2005-2008). *J. Am. Anim. Hosp. Assoc.* 49, 190– 196.

Bigge, L.A. és mtsai (2001): Correlation between coagulation profile findings and bleeding complications after ultrasound-guided biopsies: 434 cases (1993-1996). *J. Am. Anim. Hosp. Assoc.* 37, 228-233.

Center, S.A. (2007): Interpretation of liver enzymes. *Vet. Clin. North Am. Small Anim. Pract.* 37, 297– 333.

Center, S.A. és mtsai (2002): Liver glutathione concentrations in dogs and cats with naturally occurring liver disease. *Am. J. Vet. Res.* 63, 1187-1197.

Clifford, C.A. és mtsai (2004): Magnetic resonance imaging of focal splenic and hepatic lesions in the dog. *J. Vet. Intern. Med.* 18, 330– 338.

Cole, T.L. és mtsai (2002): Diagnostic comparison of needle and wedge biopsy specimens of the liver in dogs and cats. *J. Am. Vet. Med. Assoc.* 220, 1483-1490.

Ettinger, Stephen J.; Feldman, Edward C.; Cote, Etienne. *Textbook of Veterinary Internal Medicine - eBook (Kindle Location 65155)*. Elsevier Health Sciences. Kindle Edition.

Gaschen, L. (2009): Update on hepatobiliary imaging. *Vet. Clin. North Am. Small Anim. Pract.* 39, 439– 467.

Kavanagh, C. és mtsai (2011): Coagulation in hepatobiliary disease. *J. Vet. Emerg. Crit Care.* 21, 589– 604.

Kemp, S.D. és mtsai (2013): A comparison of hepatic sonographic features and histopathologic diagnosis in canine liver disease: 138 cases. *J. Vet. Intern. Med.* 27, 806– 813.

Langlois, D.K. és mtsai (2013): Acquired proximal renal tubular dysfunction in 9 Labrador retrievers with copper-associated hepatitis (2006-2012). *J. Vet. Intern. Med.* 27, 491– 499.

Larson, M.M. (2016): Ultrasound imaging of the hepatobiliary system and pancreas. *Vet. Clin. North Am. Small Anim. Pract.* 46, 453-80.

Levien, A.S. és mtsai (2014): Assessment of the efficacy and potential complications of transjugular liver biopsy in canine cadavers. *J. Vet. Intern. Med.* 28, 338– 345.

Marolf, A.J. (2016): Computed tomography and MRI of the hepatobiliary system and pancreas. *Vet. Clin. North Am. Small Anim. Pract.* 46, 481-497.

- Marolf, A.J. (2017): Diagnostic imaging of the hepatobiliary system: An update. *Vet. Clin. North Am. Small Anim. Pract.* 47, 555-568.
- Prins, M.J. és mtsai (2010): Coagulation disorders in dogs with hepatic disease. *Vet. J.* 185, 163-168.
- Roth, L. (2001): Comparison of liver cytology and biopsy diagnoses in dogs and cats: 56 cases. *Vet. Clin. Pathol.* 30, 35– 38.
- Rothuizen, J. és Twedt, D.C. (2009): Liver biopsy techniques. *Vet. Clin. North Am. Small Anim. Pract.* 39, 469– 480.
- Sterczer, Á., Reiczigel, J., Psáder, R., Pápa, K., Vörös, K., Pap, Á. (2012): Chologogue-induced gallbladder emptying in the diagnosis of canine biliary obstruction. *Acta. Vet. Hung.*, 60, 199-209.
- Sterczer, Á., Szénási, G., Vörös, K., Pap, A. (2000): Effects of lorglumide and atropine on MgSO(4)-induced gallbladder emptying in conscious dogs. *Res. Vet. Sci.* 69, 129-33.
- Sterczer, Á., Vörös, K., Karsai, F. (1996): Effect of cholagogues on the volume of the gallbladder of dogs. *Res. Vet. Sci.* 60, 44-47.
- Szatómári, V., Harkányi, Z., Vörös, K. (2003): A review of nonconventional ultrasound techniques and contrast-enhanced ultrasonography of noncardiac canine disorders. *Vet. Radiol. Ultrasound* 44, 380-391.
- Toulza, O. és mtsai (2006): Evaluation of plasma protein C activity for detection of hepatobiliary disease and portosystemic shunting in dogs. *J. Am. Vet. Med. Assoc.* 229, 1761– 1771.
- Ullal, T. és mtsai (2018): The evaluation of cyclosporine in the treatment of chronic hepatitis in dogs. In *Proceedings of ACVIM 2018. June 13-16 Seattle, Washington.*
<https://www.vin.com/acvim/2018> Utolsó hozzáférés: 2019. szept. 12.
- Ullal, T. és mtsai (2018): The evaluation of cyclosporine in the treatment of chronic hepatitis in dogs. *J. Vet. Intern. Med.* doi: 10.1111/jvim.15591. [Epub ahead of print]
- Wang, K.Y. és mtsai (2004): Accuracy of ultrasound-guided fine-needle aspiration of the liver and cytologic findings in dogs and cats: 97 cases (1990-2000). *J. Am. Vet. Med. Assoc.* 224, 75– 78.
- Webster, C.R.L. és mtsai (2019): ACVIM consensus statement on the diagnosis and treatment of chronic hepatitis in dogs. *J. Vet. Intern. Med.* 33, 1173-1200.

MÁJBETEGSÉGEK

- Azumi, N. (1982): Copper and liver injury— experimental studies on the dogs with biliary obstruction and copper loading. *Hokkaido Igaku Zasshi.* 57, 331-349.

- Balboni, A. és mtsai (2014): Investigation of the presence of canine adenovirus (CAV) in owned dogs in Northern Italy. *Res. Vet. Sci.* 97, 631– 636.
- Balkman, C.E. és mtsai (2003): Evaluation of urine sulfated and nonsulfated bile acids as a diagnostic test for liver disease in dogs. *J. Am. Vet. Med. Assoc.* 222, 1368-1375.
- Bárány, Z., Sterczler, Á., Jocsák, G., Frenyó, V.L., Kiss, D.S.: A hepaticus encephalopathia kóroktana, patogenezisének újabb aspektusai. *Magy. Állatorvosok Lapja* 139, 157-168.
- Basu, A.K. és mtsai (2014): A review of the cat liver fluke *Platynosomum fastosum* Kossack, 1910 (Trematoda: Dicrocoeliidae). *Vet. Parasitol.* 200, 1-7.
- Blanchard, G. és mtsai (2002): Dietary L-carnitine supplementation in obese cats alters carnitin metabolism and decreases ketosis during fasting and induced hepatic lipidosis. *J. Nutr.* 132, 204-210.
- Center, S.A. és mtsai (2000): Proteins invoked by vitamin K absence and clotting times in clinically ill cats. *J. Vet. Intern. Med.* 14, 292-297.
- Chung, J.Y. és mtsai (2012): Zonisamide monotherapy for idiopathic epilepsy in dogs. *N. Z. Vet. J.* 60, 357-359.
- Dirksen, K és Fieten, H. (2017): Canine copper-associated hepatitis. *Vet. Clin. North Am. Small Anim. Pract.* 47, 631-644.
- Dyggve, H. és mtsai (2011): Association of Doberman hepatitis to canine major histocompatibility complex II. *Tissue Antigens.* 77, 30– 35.
- Dyggve, H. és mtsai (2017): Antihistone autoantibodies in Dobermans with hepatitis. *J. Vet. Intern. Med.* 31, 1717-1723.
- Feeney, D.A. és mtsai (2008): Statistical relevance of ultrasonographic criteria in the assessment of diffuse liver disease in dogs and cats. *Am. J. Vet. Res.* 69, 212– 221.
- Fieten, H. és mtsai (2012): Association of dietary copper and zinc levels with hepatic copper and zinc concentrations in Labrador Retrievers. *J. Vet. Intern. Med.* 26, 1274-1280.
- Fieten, H. és mtsai (2014): Nutritional management of inherited copper-associated hepatitis in Labrador Retrievers. *Vet. J.* 199, 429-433.
- Gerritzen-Bruning M.J., és mtsai (2006): Diagnostic value of fasting plasma ammonia and bile acid concentration in the identification of portosystemic shunting in dogs. *J. Vet. Intern. Med.* 20:13-19.
- Gow, A. és mtsai (2010). Whole blood manganese concentrations I dogs with congenital portosystemic shunts. *J. Vet. Intern. Med.* 24, 90-96.
- Gow, A., és mtsai (2012): Dogs with congenital porto-systemic shunting (cPSS) and hepatic encephalopathy have higher serum concentrations of C-reactive protein than asymptomatic dogs with cPSS. *Metabolic Brain Dis.* 27, 227-229.

- Haney, D.R. és mtsai (2006): Severe cholestatic liver disease secondary to liver fluke (*Platynosomum concinnum*) infection in three cats. *J. Am. Anim. Hosp. Assoc.* 42, 234-237.
- Haywood, S. és mtsai (2016): Copper toxicosis in non-COMMD1 Bedlington terriers is associated with metal transport gene ABCA12. *J. Trace Elem. Med. Biol.* 35, 83-89.
- Hoffman, G. és mtsai (2009): Dietary management of hepatic copper accumulation in Labrador Retrievers. *J. Vet. Intern. Med.* 23, 957-963.
- Honeckman, A. (2003): Current concepts in the treatment of canine chronic hepatitis. *Clin. Tech. Small Anim. Pract.* 18, 239– 244.
- Hyun, C. és mtsai (2004): Evaluation of haplotypes associated with copper toxicosis in Bedlington Terriers in Australia. *Am. J. Vet. Res.* 65, 1573– 1579.
- Jerzsele, Á., Karancsi, Z., Pászti-Gere, E., Sterczer, Á., Bersényi, A., Fodor, K., Szabó, D., Vajdovich, P. (2018): Effects of p.o. administered xylitol in cats. *J. Vet. Pharmacol. Ther.* 41, 409-414.
- Johnston, A.N. és mtsai (2013): Hepatic copper concentrations in Labrador Retrievers with and without chronic hepatitis: 72 cases (1980-2010). *J. Am. Vet. Med. Assoc.* 242, 372-380.
- Lehel, J. és Vetter, J. (2002): Gyakoribb növényi eredetű mérgezések a kisállatpraxisban. *Magy. Állatorvosok Lapja* 124, 597-606.
- Lidbury, J.A. és mtsai (2016): Hepatic encephalopathy in dogs and cats. *J. Vet. Emerg. Crit. Care.* 26, 471–487.
- Mandigers és mtsai (2004): Chronic hepatitis in Doberman pinschers. A review. *Vet. Q.* 26, 98-106.
- Meertens, N.M. és mtsai (2005): Copper-associated chronic hepatitis and cirrhosis in a European Shorthair cat. *Vet. Pathol.* 42, 97-100.
- Otte, C.M. és mtsai (2013): Retrospective comparison of prednisolone and ursodeoxycholic acid for the treatment of feline lymphocytic cholangitis. *Vet. J.* 195, 205– 209.
- Piscitelli, C.M. és mtsai (2010): Xylitol toxicity in dogs. *Compend. Contin. Educ. Vet.* 32, E1-4.
- Podell, M. és mtsai (2016): 2015 ACVIM small animal consensus statement on seizure management in dogs. *J. Vet. Intern. Med.* 30, 477-490.
- Puschner, B. és mtsai (2007): Diagnosis of Amanita toxicosis in a dog with acute hepatic necrosis. *J. Vet. Diagn. Invest.* 19, 312– 317.
- Rothuizen, J. (2009): Important clinical syndromes associated with liver disease. *Vet. Clin. North Am. Small Anim. Pract.* 39, 419-437.
- Ruland, K. és mtsai (2010): Sensitivity and specificity of fasting ammonia and serum bile acids in the diagnosis of portosystemic shunts in dogs and cats. *Vet. Clin. Pathol.* 39, 57-64.

Steiner, J.M. és mtsai (2003): Urine sulfated and nonsulfated bile acids as a diagnostic test for liver disease in cats. *J. Vet. Intern. Med.* 17, 605.

Sterczer, Á. (1998): New diagnostic methods for liver diseases of dogs. PhD-értekezés, Állatorvostudományi Egyetem, Állatorvostudományi Doktori Iskola, Budapest.

Straten van, G. és mtsai (2015): Diagnostic value of the rectal ammonia tolerance test, fasting plasma ammonia and fasting plasma bile acids for canine portosystemic shunting. *Vet. J.* 204, 282-286.

Szatmári, V. és mtsai (2004): Ultrasonographic findings in dogs with hyperammonemia: 90 cases (2000-2002). *J. Am. Vet. Med. Assoc.* 224, 717-727.

Tivers, M.S. és mtsai (2014): Hyperammonemia and systemic inflammatory response syndrome predicts presence of hepatic encephalopathy in dogs with congenital portosystemic shunts. *PLoS One.* 9, e82303.

Trainor, D. és mtsai (2003): Urine sulfated and nonsulfated bile acids as a diagnostic test for liver disease in cats. *J. Vet. Intern. Med.* 17, 145-153.

Twedt, D.C. és mtsai (2014): Evaluation of FISH for the detection of bacteria in feline inflammatory liver disease. *J. Feline Med. Surg.* 16, 109-117.

Valtolina, C. és Favier, R. (2016): Feline hepatic lipidosis. *Vet. Clin. North Am. Small Anim. Pract.* 47, 683-702.

Von Klopmann, T. és mtsai (2007): Prospective study of zonisamide therapy for refractory idiopathic epilepsy in dogs. *J. Small Anim. Pract.* 48, 134-138.

Wagner, K.A. és mtsai (2007): Bacterial culture results from liver, gallbladder, or bile in 248 dogs and cats evaluated for hepatobiliary disease: 1998-2003. *J. Vet. Intern. Med.* 21, 417-424.

Zwingenberger, A.L. és mtsai (2014): Correlation between liver volume, portal vascular anatomy, and hepatic perfusion in dogs with congenital portosystemic shunt before and after placement of ameroid constrictors. *Vet. Surg.* 43, 926-34.

A MÁJ PORTALIS ÉRRENDSZERÉNEK BETEGSÉGEI, EGYÉB MÁJBETEGSÉGEK

Bhandal, J. és mtsai (2008): Spontaneous left medial liver lobe torsion and left lateral lobe infarction in a rottweiler. *Can. Vet. J.* 49, 1002-1004.

Bruehschwein, A. és mtsai (2010): Contrast-enhanced magnetic resonance angiography for diagnosis of portosystemic shunts in 10 dogs. *Vet. Radiol. Ultrasound.* 51, 116–121.

Buob, S. és mtsai (2011): Portal hypertension: Pathophysiology, diagnosis, and treatment. *J. Vet. Intern. Med.* 25, 169– 186.

- Echandi, R.L. és mtsai (2007): Comparison of transsplenic multidetector CT portography to multidetector CT-angiography in normal dogs. *Vet. Radiol. Ultrasound*. 48, 38-44.
- Frankel, D. és mtsai (2006): Evaluation of cellophane banding with and without intraoperative attenuation for treatment of congenital extrahepatic portosystemic shunts in dogs. *J. Am. Vet. Med. Assoc.* 228, 1355– 1360.
- Fukushima, K. és mtsai (2014): Computed tomographic morphology and clinical features of extrahepatic portosystemic shunts in 172 dogs in Japan. *Vet. J.* 199, 376-381.
- Gómez-Ochoa, P. és mtsai (2011): Use of transsplenic injection of agitated saline and heparinized blood for the ultrasonographic diagnosis of macroscopic portosystemic shunts in dogs. *Vet. Radiol. Ultrasound*. 52, 103-106.
- Goode, K. és mtsai (2019): Evaluation of hepatic tumor portal perfusion using mesenteric angiography: A pilot study in 5 dogs. *J. Vet. Intern. Med.* 33, 776-782.
- Görlinger, S. és mtsai (2003): Congenital dilatation of the bile ducts (Caroli's disease) in young dogs. *J. Vet. Intern. Med.* 17, 28-32.
- Greenhalg, S., és mtsai (2010). Comparison of survival after surgical or medical treatment in dogs with a congenital portosystemic shunt. *J. Am. Vet. Med. Assoc.* 236, 1215-1220.
- Greenhalg, S., és mtsai (2014). Long term survival and quality of life in dogs with clinical signs associated with a congenital portosystemic shunt after surgical or medical treatment. *J. Am. Vet. Med. Assoc.* 245, 527-533.
- Holt, D.E. (2002): Cerebrospinal fluid glutamine, tryptophan, and tryptophan metabolite concentrations in dogs with portosystemic shunts. *Am. J. Vet. Res.* 63, 1167-1171.
- Hostnik, E.T. (2019): Computed tomography of lobular dissecting hepatitis in a young Golden Retriever. *J. Am. Anim. Hosp. Assoc.* 55.(2) e55203
- Hunt, G.B. és mtsai (2004): Outcomes of cellophane banding for congenital portosystemic shunts in 106 dogs and 5 cats. *Vet. Surg.* 33, 25-31.
- Khoshnegah, J. és Movassaghi, A.R. (2010): A very severe case of feline amyloidosis with spontaneous hepatic rupture and chronic renal failure. *Comp. Clin. Pathol.* 19, 519– 522.
- Lamb, C.R. és Daniel, G.B. (2002): Diagnostic imaging of dogs with suspected portosystemic shunting. *Compend. Contin. Educ. Pract. Vet.* 24, 626.
- Linden, D.S. és mtsai (2019): Outcomes and prognostic variables associated with central division hepatic lobectomies: 61 dogs. *Vet. Surg.* 48, 319-314.
- Liptak, J.M. (2004): Massive hepatocellular carcinoma in dogs: 48 cases (1992-2002). *J. Am. Vet. Med. Assoc.* 225, 1225-1230.
- Loeven, K.O. (1994): Spontaneous hepatic rupture secondary to amyloidosis in a Chinese Shar Pei. *J. Am. Anim. Hosp. Assoc.* 30, 577– 579.

- Mai, W. és Weisse, C. (2011): Contrast-enhanced portal magnetic resonance angiography in dogs with suspected congenital portal vascular anomalies. *Vet. Radiol. Ultrasound*. 52, 284–288.
- Morandi, F. és mtsai (2005): Use of $^{99m}\text{TcO}_4(-)$ trans-splenic portal scintigraphy for diagnosis of portosystemic shunts in 28 dogs. *Vet. Radiol. Ultrasound* 46. 153-161.
- Nazarali, A. és mtsai (2014): Chronic liver lobe torsion in a cat. *J. Am. Anim. Hosp. Assoc.* 50, 119-123.
- Nelson, N.C. és Nelson, L.L. (2011): Anatomy of extrahepatic portosystemic shunts in dogs as determined by computed tomography angiography. *Vet. Radiol. Ultrasound*. 52, 498-506.
- Ruland, K. és mtsai (2010): Sensitivity and specificity of fasting ammonia and serum bile acids in the diagnosis of portosystemic shunts in dogs and cats. *Vet. Clin. Pathol.* 39, 57-64.
- Schwartz, S.G. (2006): Liver lobe torsion in dogs: 13 cases (1995-2004). *J. Am. Vet. Med. Assoc.* 228, 242-247.
- Schwartz, S.G. és mtsai (2006): Liver lobe torsion in dogs: 13 cases (1995-2004). *J. Am. Vet. Med. Assoc.* 228, 242-7.
- Segev, G. és mtsai (2012): Renal amyloidosis in dogs: a retrospective study of 91 cases with comparison of the disease between Shar-Pei and non-Shar-Pei dogs. *J. Vet. Intern. Med.* 26, 259– 268.
- Szatmári, V. és mtsai (2003): Intraoperative ultrasonography of the portal vein during attenuation of intrahepatic portocaval shunts in dogs. *J. Am. Vet. Med. Assoc.* 222, 1086-1092.
- Szatmári, V. és mtsai (2004a): Standard planes for ultrasonographic examination of the portal system in dogs. *J. Am. Vet. Med. Assoc.* 224, 713-716, 698-699.
- Szatmári, V. és mtsai (2004b): Ultrasonographic evaluation of partially attenuated congenital extrahepatic portosystemic shunts in 14 dogs. *Vet. Rec.* 155, 448-456.
- Tobias, K.M. (2005): Portosystemic shunts.
<https://vetmed.illinois.edu/wp-content/uploads/2015/09/54.-Portosystemic-Shunts.pdf> Utolsó hozzáférés: 2019. szept. 12.
- Tobias, K.M. és mtsai (2003): Association of breed with the diagnosis of congenital portosystemic shunts in dogs: 2,400 cases (1980-2002). *J. Am. Vet. Med. Assoc.* 223, 1636–1639.
- Toulsa, O. és mtsai (2006): Evaluation of plasma protein C activity for detection of hepatobiliary disease and portosystemic shunting in dogs. *J. Am. Vet. Med. Assoc.* 229, 1761-1771.
- Vizi Zs., Németh, T., Sterczler, Á., (2013): Comparison of the survival and quality of life in dogs with medically versus surgically treated congenital portosystemic shunts in Hungary. *Tierärztl. Prax. Ausg. K. Kleintiere Heimtiere.* 2013. 41. Paper A30.

White, R.N. és mtsai (2017): New perspectives on the development of extrahepatic portosystemic shunts. *J. Small Anim. Pract.* 58, 669-677.

Zwingenberger, A. és mtsai (2005): Helical computed tomographic angiography of canine portosystemic shunts. *Vet. Radiol. Ultrasound.* 46, 27.

AZ EPEHÓLYAG ÉS AZ EPEUTAK BETEGSÉGEI

Bennett, S.L. és mtsai (2007): Gallbladder mucocele and concurrent hepatic lipidosis in a cat. *Aust. Vet. J.* 85, 397–400.

Besso, J.G. és mtsai (2000): Ultrasonographic appearance and clinical findings in 14 dogs with gallbladder mucocele. *Vet. Radiol. Ultrasound.* 41, 261–271.

Choi, J. és mtsai (2014): Comparison between ultrasonographic and clinical findings in 43 dogs with gallbladder mucoceles. *Vet. Radiol. Ultrasound.* 55, 202–207.

Kanemoto, H. és mtsai (2017): Intrahepatic cholelithiasis in dogs and cats: A case series. *Can. Vet. J.* 58, 971–973.

Sterczer, Á., Reiczigel, J., Psáder, R., Pápa, K., Vörös, K., Pap, Á. (2012): Chologogue-induced gallbladder emptying in the diagnosis of canine biliary obstruction. *Acta Vet Hung.* 60, 199-209.

Sterczer, Á., Szénási, G., Vörös, K., Pap, Á. (2000): Effects of lorglumide and atropine on MGSO₄-induced gallbladder emptying in conscious dogs. *Res. Vet. Sci.*, 69, 129-133.

Sterczer, Á., Vörös, K., Karsai, F. (1996): Effect of cholagogues on the volume of the gallbladder of dogs. *Res. Vet. Sci.* 60, 44-47.

Vörös, K., Sterczer, Á., Manczur, F., Gaál, T. (2002): Percutaneous ultrasound-guided cholecystocentesis in dogs. *Acta Vet. Hung.* 50, 385-393.

Vörös, K., Németh, T., Vrabély, T., Manczur, F., Tóth, J., Magdus, M., Perge, E. (2001): Ultrasonography and surgery of canine biliary diseases. *Acta Vet. Hung.* 49, 141-154.

Vörös, K., Németh, T., Vrabély, T., Manczur, F., Tóth, J., Papp, L., Biksi, I., Perge, E. (2002): A kutyák elzáródásos epehólyag- és epeúti betegségeinek kórjelzése és gyógykezelése. *Magy. Állatorvosok Lapja* 124, 479-490.