

## **7. FEJEZET. A VESE ÉS A VIZELETTELVEZETŐ SZERVEK BETEGSÉGEI**

### **ÁLTALÁNOS ISMERETEK**

Finco, D.R. (2005): Measurement of glomerular filtration rate via urinary clearance of inulin and plasma clearance of technetium Tc 99m pentetate and exogenous creatinine in dogs. *Am. J. Vet. Res.* 66, 1046-1055.

Ghys, L. és mtsai (2014): Cystatin C: a new renal marker and its potential use in small animal medicine. *J. Vet. Intern. Med.* 28, 1152-1164.

McAloney, C.A. és mtsai (2018): Evaluation of the diagnostic utility of cytologic examination of renal fine-needle aspirates from dogs and the use of ultrasonographic features to inform cytologic diagnosis. *J. Am. Vet. Med. Assoc.* 252, 1247–1256.

Nabity, M.B.: Traditional renal biomarkers and new approaches to diagnostics. *Toxicol. Pathol.* 46, 999-1001.

Pelander, L. és mtsai (2019): Comparison of the diagnostic value of symmetric dimethylarginine, cystatin C, and creatinine for detection of decreased glomerular filtration rate in dogs. *J. Vet. Intern. Med.* 33, 630-639.

Relford, R. és mtsai (2016): Symmetric dimethylarginine: improving the diagnosis and staging of chronic kidney disease in small animals. *Vet. Clin. North Am. Small Anim. Pract.* 46, 941-960.

Rossi, G. és mtsai (2012): Evaluation of factors that affect analytic variability of urine protein-to-creatinine ratio determination in dogs. *Am. J. Vet. Res.* 73, 779-788.

Rossi, G. és mtsai (2015): The effect of inter-laboratory variability on the protein:creatinine (UPC) ratio in canine urine. *Vet. J.* 204, 66-72.

Watson, A.D. és mtsai (2002): Plasma exogenous creatinine clearance test in dogs: comparison with other methods and proposed limited sampling strategy. *J. Vet. Intern. Med.* 16, 22-33.

### **A VESÉK MEGBETEGEDÉSEI ÉS A VESEBETEGSÉGEK KLINIKAI FELOSZTÁSA**

Brown, S.A. és mtsai (2013): Consensus recommendations for standard therapy of glomerular disease in dogs. *J. Vet. Intern. Med.* 27, 27–43.

Burgess, K.E. és DeRegis C.J. (2019): Urologic Oncology. *Vet. Clin. North Am. Small Anim. Pract.* 49, 311-323.

Cianciolo, R.E. és mtsai (2016): World Small Animal Veterinary Association Renal Pathology Initiative: Classification of glomerular diseases in dogs. *Vet. Pathol.* 53, 113-135.

Cowgill, L.D. és Guillaumin, J. (2013): Extracorporeal renal replacement therapy and blood purification in critical care. *J. Vet. Emerg. Crit. Care* 23, 194-204.

Harjes, L.M. és mtsai (2017): Fibroblast growth factor-23 concentration in dogs with chronic kidney disease. *J. Vet. Intern. Med.* 31, 784–790.

Hartmann, K. és mtsai (2013): *Leptospira* species infection in cats: ABCD guidelines on prevention and management. *J. Feline Med. Surg.* 15, 576-581.

Hokamp, J.A. és mtsai (2018): Correlation of electrophoretic urine protein banding patterns with severity of renal damage in dogs with proteinuric chronic kidney disease. *Vet. Clin. Pathol.* 47, 425-434.

Jüppner, H. (2011): Phosphate and FGF-23. *Kidney Int. Suppl.* 79, 24–27.

Klopfleisch, R. és mtsai (2010): An emerging pulmonary haemorrhagic syndrome in dogs: similar to the human leptospiral pulmonary haemorrhagic syndrome? *Vet. Med. Int.* 2010:928541.

Knöpfler, S. és mtsai (2017): Evaluation of clinical, laboratory, imaging findings and outcome in 99 dogs with leptospirosis. *J. Small Anim. Pract.* 58, 582-588.

McAloney, C.A. és mtsai (2018a): Evaluation of the diagnostic utility of cytologic examination of renal fine-needle aspirates from dogs and the use of ultrasonographic features to inform cytologic diagnosis. *J. Am. Vet. Med. Assoc.* 252, 1247-1256.

McAloney, C.A. és mtsai (2018b): Diagnostic utility of renal fine-needle aspirate cytology and ultrasound in the cat. *J. Feline Med. Surg.* 20, 544-553.

Schuller, S. és mtsai (2015): European consensus statement on leptospirosis in dogs and cats. *J. Small Anim. Pract.* 56, 159-179.

Song, S.H. és mtsai (2014): Positron emission tomography/computed tomography imaging features of renal cell carcinoma and pulmonary metastases in a dog. *Can. Vet. J.* 55, 466-470.

Szatmári, V., Ósi, Z., Manczur, F. (2001): Ultrasound-guided percutaneous drainage for treatment of pyonephrosis in two dogs. *J. Am. Vet. Med. Assoc.* 218, 1796–1799.

## **A VIZELETÉLVEZETŐ SZERVEK BETEGSÉGEI**

Adams, L. és mtsai (2008): Use of laser lithotripsy for fragmentation of uroliths in dogs: 73 cases (2005-2006). *J. Am. Vet. Med. Assoc.* 11, 1680-1687.

Albasan, H. és mtsai (2012): Risk factors for urate uroliths in cats. *J. Am. Vet. Med. Assoc.* 240, 842–847.

Bannasch, D. és mtsai (2008): Mutations in the SLC2A9 gene cause hyperuricosuria and hyperuricemia in the dog. *PLoS Genet.* 4, e1000246.

- Bende B. és Németh T. (2004): High prevalence of urate urolithiasis in the Russian black terrier. *Vet. Rec.* 155, 239-240.
- Bende, B. és Németh, T. (2015): Macskák húgykövességének epidemiológiai vizsgálata Magyarországon 2006 és 2014 között (480 eset). *Magy. Állatorvosok Lapja* 137, 305-313.
- Bende, B., Kovács, K. B., Solymosi, N., Németh T. (2015): Characteristics of urolithiasis in the dog population of Hungary from 2001 to 2012. *Acta Vet. Hung.* 63, 323-336.
- Berent, A.C. és mtsai (2008): Use of cystoscopic-guided laser ablation for treatment of intramural ureteral ectopia in male dogs: four cases (2006-2007). *J. Am. Vet. Med. Assoc.* 232, 1026-1034.
- Berent, A.C. és mtsai (2018): Use of a subcutaneous ureteral bypass device for treatment of benign ureteral obstruction in cats: 174 ureters in 134 cats (2009-2015). *J. Am. Vet. Med. Assoc.* 253, 1309-1327.
- Blackburn, A.L. és mtsai (2013): Evaluation of outcome following urethral stent placement for the treatment of obstructive carcinoma of the urethra in dogs: 42 cases (2004-2008). *J. Am. Vet. Med. Assoc.* 242, 59-68.
- Brons, A.K. és mtsai (2013): SLC3A1 and SLC7A9 mutations in autosomal recessive or dominant canine cystinuria: A new classification system. *J. Vet. Intern. Med.* 27, 1400-1408.
- Cannon, A.B. és mtsai (2007): Evaluation of trends in urolith composition in cats: 5,230 cases (1985-2004). *J. Am. Vet. Med. Assoc.*, 231, 570–576.
- Cunto, M. és mtsai (2019): Clinical approach to prostatic diseases in the dog. *Reprod. Domest. Anim.* 54, 816-822.
- Del Angel-Caraza, J. és mtsai (2010): Composition of lower urinary tract stones in canines in Mexico City. *Urol. Res.* 38, 201-204.
- Gallagher, A. (2018): Interventional radiology and interventional endoscopy in treatment of nephroureteral disease in the dog and cat. *Vet. Clin. North Am. Small Anim. Pract.* 48, 843-862.
- Ghantous, S.N. és Crawford J. (2006): Double ureters with ureteral ectopia in a domestic shorthair cat. *J. Am. Anim. Hosp. Assoc.* 42, 462-456.
- Heilmann, R.M. (2016): Minimally invasive interventional techniques involving the urogenital tract in dogs and cats. *Tierärztl. Prax. Ausg. K. Kleintiere Heimtiere.* 44, 105-115.
- Henthorn, P.S. és mtsai (2000): Canine cystinuria: polymorphism in the canine SLC3A1 gene and identification of a nonsense mutation in cystinuric Newfoundland dogs. *Hum. Gen.* 107, 295–303.
- Lekcharoensuk, C. és mtsai (2000): Patient and environmental factors associated with calcium oxalate urolithiasis in dogs. *J. Am. Vet. Med. Assoc.* 217, 515–519.
- Low, W.W. és mtsai (2010): Evaluation of trends in urolith composition and characteristics of dogs with urolithiasis: 25,499 cases (1985-2006). *J. Am. Vet. Med. Assoc.* 236, 193–200.

Lulich J.P. és mtsai (2009): Efficacy and safety of laser lithotripsy in fragmentation of urocalculi and urethroliths for removal in dogs. *J. Am. Vet. Med. Assoc.* 10, 1279-1285.

Lulich, J.P. és mtsai (2013): Efficacy of two commercially available, low-magnesium, urine-acidifying dry foods for the dissolution of struvite urolith in cats. *J. Am. Vet. Med. Assoc.* 8, 1147-1153.

Lulich, J.P. és mtsai (2013): Recent shifts in the global proportions of canine uroliths. *Vet. Rec.* 172, 363-369.

Lulich, J.P. és mtsai (2016): ACVIM Small animal consensus recommendation on the treatment and prevention of uroliths in dogs and cats. *J. Vet. Intern. Med.* 30, 1564-1574.

Macrì, F. és mtsai (2018): Quantitative evaluation of canine urinary bladder transitional cell carcinoma using contrast-enhanced ultrasonography. *BMC Vet. Res.* 14, 84.

McMillan, S.K. és mtsai (2011): Antitumor effects of deracoxib treatment in 26 dogs with transitional cell carcinoma of the urinary bladder. *J. Am. Vet. Med. Assoc.* 239, 1084-1089.

Mizukami, K. és mtsai (2015): Feline cystinuria caused by a missense mutation in the SLC3A1 gene. *J. Vet. Intern. Med.* 29, 120-125.

Mizukami, K. és mtsai (2016): Cystinuria associated with different SLC7A9 gene variants in the cat. *PLoS One.* 11, e0159247.

Naughton, J.F. és mtsai (2012): Accuracy of three-dimensional and two-dimensional ultrasonography for measurement of tumor volume in dogs with transitional cell carcinoma of the urinary bladder. *Am. J. Vet. Res.* 73, 1919-1924.

Osborne, C.A. és mtsai (2009): Analysis of 451,891 canine uroliths, feline uroliths, and feline urethral plugs from 1981 to 2007: perspectives from the Minnesota urolith center. *Vet. Clin. North Am. Small Anim. Pract.* 39, 183–197.

Runge, J.J. és mtsai (2011): Transvesicular percutaneous cystolithotomy for the retrieval of cystic and urethral calculi in dogs and cats: 27 cases (2006-2008). *J. Am. Vet. Med. Assoc.* 3, 344-349.

Schenk, F. és mtsai (2010): Silica containing uroliths in dogs from Switzerland: A remarkable increase in frequency after 2006. In: *Proceedings of the 20th ECVIM-CA Congress, Toulouse, France*, 303.

Tobias, K.M. és mtsai (2003): Association of breed with the diagnosis of congenital portosystemic shunts in dogs: 2,400 cases (1980-2002). *J. Am. Vet. Med. Assoc.* 223, 1636–1639.

## **A VIZELETÜRÍTÉS ZAVARAI**

Applegate, R. és mtsai (2018): Urethral sphincter mechanism incompetence in dogs: an update. *J. Am. Anim. Hosp. Assoc.* 54, 22-29.

Berent, A.C. és mtsai (2008): Use of cystoscopic-guided laser ablation for treatment of intramural ureteral ectopia in male dogs: four cases (2006-2007). *J. Am. Vet. Med. Assoc.* 232, 1026-1034.

Berent, A.C. és mtsai (2012): Evaluation of cystoscopic-guided laser ablation of intramural ectopic ureters in female dogs. *J. Am. Vet. Med. Assoc.* 240, 716-725.

Cunto, M. és mtsai (2019): Clinical approach to prostatic diseases in the dog. *Reprod. Domest. Anim.* 54, 816-822.

Ghantous, S.N. és Crawford, J. (2006): Double ureters with ureteral ectopia in a domestic shorthair cat. *J. Am. Anim. Hosp. Assoc.* 42, 462-466.

Lisciandro, G.R. és Fosgate, G.T. (2017): Use of urinary bladder measurements from a point-of-care cysto-colic ultrasonographic view to estimate urinary bladder volume in dogs and cats. *J. Vet. Emerg. Crit. Care.* 27, 713-717.